Coset CFTs, non-abelian T-duality and high spin sectors

Konstadinos Sfetsos

Department of Engineering Sciences, University of Patras, GREECE

 $OP\ThetaO\Delta O\Xi H AKA\Delta HMIA KPHTH\Sigma$, 16 September 2010

Work with A.P. Polychronakos (City University of New York)

- Nucl.Phys. B840 (2010) 534-552, arXiv:1006.2386 [hep-th].
- arXiv:1008.3909 [hep-th].

Motivation

Field equations in curved stringy backgrounds

In string theory want to go beyond (super)gravity:

- ▶ \exists exact theories realizing this, particularly, coset (*G*/*H*) CFTs .
- ► When groups are non-abelian there are no isometries (generic).
- Solving the field equations is an impossible task with traditional methods, i.e. separation of variables.
- In physical applications this is precisely what is needed, i.e. propagating fluctuations, etc.

Understanding Non-abelian T-duality

Unlike abelian T-duality:

- Not well understood.
- Not likely to be an exact symmetry.
- Yet, what is it good for? Maybe for some effective description?

(ロ) (型) (E) (E) (E) (O)

Outline

- Gauged WZW models and Non-abelian T-duality:
 - Gauged WZW models and their geometry.
 - ► Non-abelian WZW T-duals and their geometry.
 - Relating non-abelian T-duality to gauged WZW models.
- Solving field equations in coset CFT backgrounds.
- Example: $SU(2) \times SU(2)/SU(2)$
- The infinitely large spin limit and the effective non-abelian T-dual.

- Example: Non-abelian T-dual of SU(2) WZW.
- Concluding remarks.

Gauged WZW models and Non-abelian T-duality

Gauged WZW models and their geometry

Let a group G, a subgroup $H \in G$. Introduce $g \in G$ and gauge fields $A_{\pm} \in \mathcal{L}(H)$.

The gauged WZW action is

$$S(g, A_{\pm}) = k \overbrace{I_0(g)}^{\mathsf{WZW}} + \frac{k}{\pi} \int \mathrm{Tr} \Big[A_- \partial_+ g g^{-1} - A_+ g^{-1} \partial_- g + A_- g A_+ g^{-1} - A_- A_+ \Big]$$

• Gauge invariance: For $\Lambda \in H$

$$g o \Lambda^{-1} g \Lambda$$
 , $A_\pm o \Lambda^{-1} (A_\pm - \partial_\pm) \Lambda$.

Gauge fix dim(H) parameters in g, leaving dim(G/H) x^{μ} 's.

• Integrating out the A_{\pm} 's and obtain

$$S = \frac{k}{\pi} \int (G_{\mu\nu} + B_{\mu\nu}) \partial_+ X^{\mu} \partial_- X^{\nu}$$
, also $\Phi = \cdots$

(ロ) (型) (E) (E) (E) (O)

▶ Isometries $G_L \times G_R$ of the WZW are generically broken.

Non-abelian WZW T-duals and their geometry

The starting action is

$$S_{\text{NonAb}}(g, v, A_{\pm}) = S(g, A_{\pm}) - i\frac{k}{\pi} \int \underbrace{\text{Tr}(vF_{+-})}_{\text{Lagrange mult.}}$$

- Gauge invariance: As before and in addition $v \to \Lambda^{-1} v \Lambda$.
- Gauge fix dim(H) parameters in g and v, leaving dim(G) variables X^M. Integrate out the A_±'s and get the σ-model.
- Properties:
 - Isometries $G_L \times G_R$ of the WZW generically broken.
 - Even if G is compact, (some) variables of its non-abelian dual appear non-compact.
 - Transformation is not invertible at the action level.
 - Drastically different than abelian T-duality.
- Is it useful? What does it describe?

Relating non-abelian T-duality to gauged WZW models

- ► Start with the gauged WZW action for $\frac{G_k \times H_\ell}{H_{k+\ell}}$ for two group elements $g \in G$, $h \in H$ and gauged field in $\mathcal{L}(H)$.
- Expand infinitesimally around the identity

$$h = \mathbb{I} + i \frac{k}{\ell} v + \mathcal{O}\left(\frac{1}{\ell^2}\right)$$

and take the limit $\ell \to \infty$.

► We get the non-abelian T-duality action, i.e. classically [KS 94]

$$\left. \frac{G_k \times H_\ell}{H_{k+\ell}} \right|_{\ell \to \infty} = \text{ dual of } G_k \text{ with respect to } H$$

Remarks:

- In the limit some variables become non-compact.
- A well defined limit can be taken on the geometric background.
- Can this be the effective background describing a consistent sector of the parent theory?

Solving field equations in coset CFT backgrounds

Consider the scalar equation (with measure $e^{-2\Phi}\sqrt{G}$) for the coset background for $G_k \times H_\ell/H_{k+\ell}$.

- We will obtain its general solution from that of the scalar equation for the WZW model for $G \times H$.
- Start with Reps of $G \times H$. The eigenstates are

 $R_{lphaeta}(g) \; r_{\mu
u}(h)$,

with eigenvalues (semiclassically, for $k, \ell \gg 1$)

$$E(R,r)=\frac{C_2(R)}{k}+\frac{C_2(r)}{\ell},$$

where the C_2 's are the Casimirs.

Under the vector H-transf they transform as

$$(R \times r) \times (\bar{R} \times \bar{r}) = (r_1 \oplus r_2 \oplus \cdots) \otimes (\bar{r}_1 \oplus \bar{r}_2 \oplus \cdots)$$
.

- We decompose $R \times r$ and its conjugate into Reps r_i of H.
- We get a singlet from all products of the form $r_i \times \bar{r}_i$.
- The coset eigenstates are



Combinations of states of the $G \times H$ WZW.

The eigenvalues get shifted as

$$E(R, r; r_i) = \frac{C_2(R)}{k} + \frac{C_2(r)}{\ell} - \frac{C_2(r_i)}{k+\ell}$$

- ► The coset background receives 1/k corrections. It becomes simple for k ≫ 1.
- ► Remarkably, the eigenstates do not depend on α' ~ 1/k, only the eigenvalues do (indicated expressions are for k ≫ 1).

Example: $SU(2) \times SU(2)/SU(2)$

The background fields

Parametrized by three compact variables

$$0\leqslant lpha_0, eta_0\leqslant 1$$
 , $|\gamma|\leqslant \sqrt{1-lpha_0^2}\sqrt{1-eta_0^2}$.

The metrics is

$$ds^{2} = \frac{k_{1} + k_{2}}{(1 - \alpha_{0}^{2})(1 - \beta_{0}^{2}) - \gamma^{2}} (\Delta_{\alpha\alpha} d\alpha_{0}^{2} + \Delta_{\beta\beta} d\beta_{0}^{2} + \Delta_{\gamma\gamma} d\gamma^{2} + 2\Delta_{\alpha\beta} d\alpha_{0} d\beta_{0} + 2\Delta_{\alpha\gamma} d\alpha_{0} d\gamma + 2\Delta_{\beta\gamma} d\beta_{0} d\gamma) .$$

The Δ 's are functions of α_0 , β_0 , γ and of $r = k_1/k_2$.

- The field $B_{\mu\nu} = 0$ and there is also a Φ .
- Background is a bit complicated, with no isometries.

Solution of the eigenvalue problem

The general state is

$$\Psi_{j_{1},j_{2}}^{j} = \sum_{m} \sum_{m_{2},n_{2}=-j_{2}}^{j_{2}} \underbrace{C_{j_{1},m-m_{2},j_{2},m_{2}}^{j,m} C_{j_{1},m-n_{2},j_{2},n_{2}}^{j,m}}_{\text{Clebsch-Gordan}} \underbrace{\frac{SU(2) \times SU(2) \text{ d-functions}}{R_{m-m_{2},m-n_{2}}^{j_{1}}(g_{1})R_{m_{2},n_{2}}^{j_{2}}(g_{2})}{\text{gauged fixed}}$$

Examples:

• For $j_2 = 0$ and thus $j_1 = j$:

$$\Psi_{j,0}^{j} = \overbrace{\sum_{m=-j}^{j} R_{m,m}^{j}(g_{1})}^{\text{Character}} = \overbrace{\underbrace{U_{2j}(\alpha_{0})}^{\text{Chebyshev}}}_{\text{2nd kind}} .$$

• For $(j_1, j_2) = (1, 1/2)$:

$$\Psi_{1,1/2}^{1/2} = 4 \alpha_0^2 - 1) \beta_0 + 4 \alpha_0 \gamma$$
, $\Psi_{1,1/2}^{3/2} = (4 \alpha_0^2 - 1) b_0 - 2 \alpha_0 \gamma$.

- ▶ With increasing *j*_{1,2} expressions become complicated.
- What about $j_1, j \gg 1$?

Large spins and the effective non-abelian T-dual High spin limit in the $G_k \times H_\ell / H_{k+\ell}$ theory?

- Let Reps in L(H) with highest weight (spin) j ≫ 1. The Reps in the tensor product with those in L(G) have also large spin.
- We may expand as

$$C_2(r) = a(r)j^2 + b(r)j + O(1)$$
.

- Similarly for $C_2(r_i)$, with j replaced by j + n (n =finite).
- Keeping the eigenenergies finite requires the correlated limit

$$\ell = rac{k}{\delta} \; j o \infty$$
 , $\delta = ext{positive real}$.

The limit of the eigenfunction is delicate. It involves the limiting behaviour of the Clebsch–Gordans.

- ▶ But, $\ell \to \infty$ is associated to the non-abelian T-dual of G_k .
- Hence:

Non-abelian T-duality provides an effective description of the high spin sector of the parent theory. *Example: Non-abelian T-dual of SU*(2) WZW

The background fields

▶ Non-abelian T-dual of the SU(2) WZW model w.r.t. SU(2)

$$ds^{2} = d\psi^{2} + \frac{\cos^{2}\psi}{x_{3}^{2}}dx_{1}^{2} + \frac{\left(x_{3}dx_{3} + (\sin\psi\cos\psi + x_{1} + \psi)dx_{1}\right)^{2}}{x_{3}^{2}\cos^{2}\psi}$$

plus a dilaton.

- A bit complicated with no isometries.
- ψ is periodic and x_1, x_3 are non-compact.
- What do the eigenfunctions and eigenenergies look like? They should effectively describe the large spin sector of the SU(2) × SU(2)/SU(2) coset.

Solution of the eigenvalue problem

We will take the limit in the states and eigenvalues of the coset.

Consider the high spin-level limit

$$j_1 = j - n$$
, $k_1 = \frac{k_2}{\delta} j$, $j_2, n = \text{finite}$, $j \gg 1$.

▶ The energy eigenvalues $E_{j_1,j_2}^j = \frac{j_1(j_1+1)}{k_1} + \frac{j_2(j_2+1)}{k_2} - \frac{j(j+1)}{k_1+k_2}$, remain finite

$$E_{j_2,n,\delta} = \lim_{j \to \infty} E_{j_1,j_2}^j = \frac{j_2(j_2+1)}{k_2} + \frac{\delta - 2n}{k_2} \,\delta$$

In the high spin limit the Clebsch–Gordan coefficients

$$\lim_{j \to \infty} C_{j-n,m-m_2,j_2,m_2}^{j,m} = d_{m_2,n}^{j_2}(\zeta) , \quad \cos \zeta = \frac{m}{j}$$

- They get associated with an auxiliary SU(2) rotation.
- Expected for a classical body given extra angular momentum.

うして ふゆう ふほう ふほう うらつ

At the end we obtain the finite sum

$$\Psi_{j_2,n,\delta}(x_1,x_3,\psi) = \lim_{j \to \infty} \Psi_{j-n,j_2}^j = \sum_{m_2=-j_2}^{j_2} \Gamma_{j_2,m_2,n,\delta}(x_3) \underbrace{\mathcal{R}_{m_2,m_2}^{j_2}(g_2)}_{\text{gauged fixed}} \right|,$$

where

$$\Gamma_{j_2,m_2,n,\delta}(x_3) = \int_0^\pi d\zeta \sin\zeta \left(d_{m_2,n}^{j_2}(\zeta)\right)^2 e^{-2i\delta v_3 \cos\zeta}$$

- ► Explicit expressions become complicated fast, as *j*₂ increases.
- Fair to say:

Solution would have never been found without using this method.

ション ふゆ アメリア メリア しょうくしゃ

Examples of states:

Define

$$v_3 = \sqrt{(x_1 + \psi)^2 + x_3^2} , \qquad \beta_0 = \sin \psi ,$$

$$\beta_1 = \frac{x_3 \cos \psi}{\sqrt{(x_1 + \psi)^2 + x_3^2}} , \qquad \beta_3 = \frac{(x_1 + \psi) \cos \psi}{\sqrt{(x_1 + \psi)^2 + x_3^2}} .$$

• For instance, for $j_2 = 1$ (and $\delta = 1$):

$$\begin{split} \Psi_{1,\pm 1} &= \frac{\beta_1^2 - 2\beta_3(\beta_3 \mp 2\beta_0 v_3)}{2v_3^2} \cos 2v_3 \\ &\quad + \frac{2\beta_3^2 - \beta_1^2 + \mp 4\beta_0\beta_3 v_3 + 4(\beta_0^2 - \beta_3^2)v_3^2}{4v_3^3} \sin 2v_3 \ , \\ \Psi_{1,0} &= \frac{2\beta_3^2 - \beta_1^2}{v_3^2} \ \cos 2v_3 + \frac{\beta_1^2 - 2\beta_3^2 + 2(1 - 2\beta_1^2)v_3^2}{2v_3^3} \ \sin 2v_3 \ . \end{split}$$

<□▶ <□▶ < □▶ < □▶ < □▶ < □ > ○ < ○

Concluding remarks

- Using group theoretical methods one may solve field equations for general G/H, for which:
 - Generically, there are no isometries.
 - Conventional techniques are not applicable.
- Non-abelian duality generates solutions that:
 - effectively describe high spin sectors.
 - Taking the limit is a delicate procedure, but nevertheless the only way to solve field equation of the T-dual background.
- Method works in other occasions with non-abelian isometries.
 For instance, when the symmetry group acts from on side, i.e. in Principal Chiral Models.
- ► Use non-compact groups leading to Minkowski signature spacetimes, i.e. SL(2, ℝ) × SL(2, ℝ) / SL(2, ℝ). Explore physical applications, i.e. in cosmology.