


Kolymbari, 13 July, 2017



Elias Kiritsis



CCTP/ITCP University of Crete APC, Paris

I. Bakas and higher spin algebras

- One of Ioannis' most important contributions to physics is the introduction and study of higher spin algebras.
- They were introduced in the context of two dimensional theories
- This came a bit after Zamolodchikov introduced the W_N algebra in 2d CFTs.

THE LARGE- N LIMIT OF EXTENDED CONFORMAL SYMMETRIES

I. BAKAS ¹

Center for Theoretical Physics, Department of Physics and Astronomy, University of Maryland, College Park, MD 20742, USA

Received 27 April 1989; revised manuscript received 14 June 1989

We study the large- N limit of the operator algebra W_N generated by primary conformal fields with integer spin $1, 2, \dots, N$. It is shown that W_∞ provides a representation of a certain infinite dimensional (sub)algebra of the area-preserving diffeomorphisms of the 2-plane (compactified or not). We also discuss certain applications of this result to quantum field theory.

- Ioannis started from the Drinfeld-Sokolov reduction $SL(N) \rightarrow W_N$ and suggested to take the limit $N \rightarrow \infty$.
- Showed that classical W_∞ algebra obtained is related to area-preserving diffeos

● In a highly prescient conclusion page he suggested that:

1. W_∞ is related to quantum theories with area preserving symmetry like membranes, hydrodynamics and fermi surfaces.
2. It is related to an “enveloping” algebra of all W_N symmetries of Zamolodchikov.
3. Suggested that the construction of W_N and W_∞ strings is possible and interesting.
4. Suggested that the $N \rightarrow \infty$ of Z_N parafermions, is a W_∞ theory that is related to area preserving diffeos.
5. Suggested that they are similar to the higher-spin theories of Fradkin and Vasiliev.

The Structure of the W_∞ Algebra

I. Bakas*


Center for Theoretical Physics, Department of Physics and Astronomy, University of Maryland, College Park, MD 20742, USA

Received November 29, 1989

Abstract. We prove rigorously that the structure constants of the leading (highest spin) linear terms in the commutation relations of the conformal chiral operator algebra W_∞ are identical to those of the $\text{Diff}_0^+ \mathbb{R}^2$ algebra generated by area preserving diffeomorphisms of the plane. Moreover, all quadratic terms of the W_N algebra are found to be absent in the limit $N \rightarrow \infty$. In particular we show that W_∞ is a central extension of $\text{Diff}_0^+ \mathbb{R}^2$ with non-trivial cocycles appearing only in the commutation relations of its Virasoro subalgebra. We also propose a representation of W_∞ in terms of a single scalar field in $2 + 1$ dimensions and discuss its significance in the context of quantum field theory.

- Fast forward to Berkeley in 1990:

Ioannis visited me, and while he came another important paper came out:



W_∞ and the Racah-Wigner algebra

C.N. Pope,* L.J. Romans[†] and X. Shen*

ABSTRACT

We examine the structure of a recently-constructed W_∞ algebra, an extension of the Virasoro algebra that describes an infinite number of fields with all conformal spins 2, 3, ..., with central terms for all spins. By examining its underlying $SL(2, R)$ structure, we are able to exhibit its relation to the algebras of $SL(2, R)$ tensor operators. Based upon this relationship, we generalise W_∞ to a one-parameter family of inequivalent Lie algebras $W_\infty(\mu)$, which for general μ requires the introduction of formally negative spins. Furthermore, we display a realisation of the $W_\infty(\mu)$ commutation relations in terms of an underlying associative product, which we denote with a lone star. This product structure shares many formal features with the Racah-Wigner algebra in angular momentum theory. We also discuss the relation between W_∞ and the symplectic algebra on a cone, which can be viewed as a co-adjoint orbit of $SL(2, R)$.

BOSONIC REALIZATION OF A UNIVERSAL W-ALGEBRA AND Z_∞ PARAFERMIONS*

Ioannis BAKAS

*Center for Theoretical Physics**, Department of Physics and Astronomy, University of Maryland,
College Park, MD 20742, USA*

Elias KIRITSIS

*Department of Physics***, University of California,
and
Theoretical Physics Group, Lawrence Berkeley Laboratory, Berkeley, CA 94720 USA*

Received 30 March 1990

We construct a field theoretic representation of the universal W-algebra proposed by Pope, Romans and Shen, using a free complex boson in two dimensions. The resulting symmetry algebra is generated by conformal fields with spin $2, 3, 4, \dots$ and has central charge $c = 2$. Highest-weight representations are also given in terms of vertex operators. Furthermore, we discuss the relation of this representation to the theory of Z_∞ parafermions.

- Our work realized the symmetry algebra as written by PRS, with a FREE complex massless boson and introduced "colored" generalizations of W_∞
- Later, Pope Romans and Shen realized $W_{1+\infty}$ in terms of a FREE Dirac Fermion.
- The way was open for realizations of W_∞ in many concrete systems.
- Almost 25 years later, Maldacena and Zhiboedov proved the reverse: only free theories have W_∞ symmetries.

BOSONIZATION OF NON-RELATIVISTIC FERMIONS AND W -INFINITY ALGEBRA

SUMIT R. DAS, AVINASH DHAR, GAUTAM MANDAL*

Tata Institute of Fundamental Research, Homi Bhabha Road, Bombay 400005, India

and

SPENTA R. WADIA^{†,‡,§}

School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540, USA

Received 18 November 1991

We discuss the bosonization of non-relativistic fermions in one-space dimension in terms of bilocal operators which are naturally related to the generators of W -infinity algebra. The resulting system is analogous to the problem of a spin in a magnetic field for the group W -infinity. The new dynamical variables turn out to be W -infinity group elements valued in the coset W -infinity/ H where H is a Cartan subalgebra. A classical action with an H gauge invariance is presented. This action is three-dimensional. It turns out to be similar to the action that describes the color degrees of freedom of a Yang-Mills particle in a fixed external field. We also discuss the relation of this action with the one recently arrived at in the Euclidean continuation of the theory using different coordinates.

Fermions in the lowest Landau level. Bosonization, W_∞ algebra, droplets, chiral bosons

Satoshi Iso ¹

Department of Physics, University of Tokyo, Bunkyo-ku, Tokyo 113, Japan

Dimitra Karabali ^{2,3} and B. Sakita ⁴

Department of Physics, City College of the City University of New York, New York, NY 10031, USA

Received 22 September 1992

We present field theoretical descriptions of massless $(2+1)$ dimensional nonrelativistic fermions in an external magnetic field, in terms of a fermionic and bosonic second quantized language. An infinite dimensional algebra, W_∞ , appears as the algebra of unitary transformations which preserve the lowest Landau level condition and the particle number. In the droplet approximation it reduces to the algebra of area-preserving diffeomorphisms, which is responsible for the existence of a universal chiral boson lagrangian independent of the electrostatic potential. We argue that the bosonic droplet approximation is the strong magnetic field limit of the fermionic theory. The relation to the $c=1$ string model is discussed.

- Boris Khesin (student of Arnold), then professor at Berkeley Math department had introduced the log of derivative cocycle to introduce central terms in algebras of pseudo-differential operators.
- We teamed up.....

The Logarithm of the Derivative Operator and Higher Spin Algebras of W_∞ Type*

Ioannis Bakas^{1} Boris Khesin^{2****} and Elias Kiritsis^{3*****}**

¹ Center for Theoretical Physics,^{***} Department of Physics and Astronomy, University of Maryland, College Park, MD 20742, USA

² Department of Mathematics, University of California, Berkeley, CA 94720, USA

³ Department of Physics,^{*****} Theoretical Physics Group, Lawrence Berkeley Laboratory, University of California, Berkeley, CA 94720, USA

Received October 22, 1991

- An important reason that made W_∞ popular was due to the expectation that the W_∞ charges would be important in the resolution of the black hole information paradox of the two dimensional black hole.

BEYOND THE LARGE N LIMIT: NON-LINEAR W_∞ AS SYMMETRY OF THE $SL(2,R)/U(1)$ COSET MODEL*

IOANNIS BAKAS⁺

*Center for Theoretical Physics
Department of Physics and Astronomy, University of Maryland
College Park, MD 20742, USA*

and

ELIAS KIRITSIS[†]

*Department of Physics, University of California and
Theoretical Physics Group, Lawrence Berkeley Laboratory
Berkeley, CA 94720, USA*

Received September 24, 1991

ABSTRACT

We show that the symmetry algebra of the $SL(2,R)_\kappa/U(1)$ coset model is a non-linear deformation of W_∞ , characterized by κ . This is a universal W -algebra which linearizes in the large κ limit and truncates to W_N for $\kappa = -N$. Using the theory of non-compact parafermions we construct a free field realization of the non-linear W_∞ in terms of two bosons with background charge. The W -characters of all unitary $SL(2,R)/U(1)$ representations are computed. Applications to the physics of 2-d black hole backgrounds are also discussed and connections with the KP approach to $c=1$ string theory are outlined.

- Since then W_∞ activity quieted down
- Vasiliev went ahead to develop to develop his higher spin theories in AdS completely unnoticed (in the 1990's)
- Things were revived in 2010.....

An AdS₃ dual for minimal model CFTs

Matthias R. Gaberdiel^{1,2,*} and Rajesh Gopakumar^{3,†}¹*School of Natural Sciences, Institute for Advanced Study, Princeton, New Jersey 08540, USA*²*Institut für Theoretische Physik, ETH Zurich, CH-8093 Zürich, Switzerland*³*Harish-Chandra Research Institute, Chhatnag Road, Jhusi, Allahabad, India 211019*

(Received 10 December 2010; published 8 March 2011)

We propose a duality between the 2d \mathcal{W}_N minimal models in the large N 't Hooft limit, and a family of higher spin theories on AdS₃. The 2d conformal field theories (CFTs) can be described as Wess-Zumino-Witten coset models, and include, for $N = 2$, the usual Virasoro unitary series. The dual bulk theory contains, in addition to the massless higher spin fields, two complex scalars (of equal mass). The mass is directly related to the 't Hooft coupling constant of the dual CFT. We give convincing evidence that the spectra of the two theories match precisely for all values of the 't Hooft coupling. We also show that the renormalization group flows in the 2d CFT agree exactly with the usual AdS/CFT prediction of the gravity theory. Our proposal is in many ways analogous to the Klebanov-Polyakov conjecture for an AdS₄ dual for the singlet sector of large N vector models.


Epilogue

- Ioannis had also several other important contributions in physics.
- Beyond physics, he was a remarkable personality: he was warm, honest, with high standards, both on moral issues and on quality.
- He will live on in our minds, as long as we live.
- But his scientific legacy will outlive all of us.

Some photos











I. Bakas and W_∞ symmetries,

Elias Kiritsis