

EPS Conference, Grenoble
23 July 2011

*The relativistic Langevin dynamics
of heavy quark diffusion from
Holography*

Elias Kiritsis



University of Crete



(APC, Paris)

Collaborators

- Umut Gursoy (CERN)
- Liuba Mazzanti (Santiago de Compostella)
- Francesco Nitti (APC, Paris 7)

Work based on ongoing work and

- U. Gursoy, E. Kiritsis, L. Mazzanti and F. Nitti,
“Langevin diffusion of heavy quarks in non-conformal holographic back-grounds”
[ArXiv:1006.3261][[hep-th]].

and previous work:

- U. Gursoy, E. Kiritsis, G. Michalogiorgakis and F. Nitti,
“ Thermal Transport and Drag Force in Improved Holographic QCD”
[ArXiv:0906.1890][[hep-ph]].

Previous work in the context of $N=4$ sYM:

- J. Casalderrey-Solana and D. Teaney, [[arXiv:hep-ph/0605199](#)].
- S. S. Gubser, [[arXiv:hep-th/0612143](#)].
- J. Casalderrey-Solana and D. Teaney, [[arXiv:hep-th/0701123](#)].
- J. de Boer, V. E. Hubeny, M. Rangamani and M. Shigemori, [[ArXiv:0812.5112](#)][hep-th].
- D. T. Son and D. Teaney, [[ArXiv:0901.2338](#)][hep-th].
- G. C. Giecold, E. Iancu and A. H. Mueller, [[ArXiv:0903.1840](#)][hep-th].

Plan of the presentation

- Introduction
- Langevin Dynamics and Brownian Motion
- Holographic computation of the Langevin diffusion.
- Relevance for RHIC and LHC
- Outlook

Introduction

- An important class of probes in Heavy ion collisions are heavy quarks.
- They are relatively easily identifiable from end products.
- They can provide useful “localized” information about the quark-gluon fireball in a heavy-ion collision.
- They have not been so prominent in the initial phases of RHIC due to energy availability.
- They are becoming more prominent and they are expected to play an important role at LHC, complementing the collective observables.

Brownian Motion and Langevin Dynamics

- Heavy particles moving inside a thermal bath undergo Brownian motion: once in a while they collide with fluid particles and suddenly change path.
- This phenomenon has an elegant description in terms of the (local) **Langevin equation** which in its simplest form is

$$\frac{dp^i(t)}{dt} = -\eta_D^{ij} p^j(t) + \xi^i(t) \quad , \quad \langle \xi^i(t) \rangle = 0 \quad , \quad \langle \xi^i(t) \xi^j(t') \rangle = 2\kappa^{ij} \delta(t - t')$$

η_D^{ij} is an average “viscous” (dissipative) force

κ^{ij} is the diffusion coefficients.

- Physically both of them have a common origin: the interactions of the heavy probe with the heat-bath.
- The first describes the averaged out (smooth) motion, while the second the (stochastic) fluctuations around the average motion.
- The Langevin equation is a stochastic equation and as such makes sense only in a concrete (time) discretized form.

The generalized Langevin equation (with memory)

- For our purposes a more general analysis is necessary. We consider the coupling of the coordinates of the probe with the bath degrees of freedom

$$S_{int} = \int d\tau \vec{X}(\tau) \cdot \vec{\mathcal{F}}$$

where \mathcal{F}^i is the force from the heat-bath.

- The **generalized Langevin equation** in general has memory and reads

$$\dot{P}^i + \int_0^\infty dt' \gamma^{ij}(t') \dot{X}^j(t-t') = \xi^i(t)$$

where

$$\dot{\gamma}^{ij}(t) = G_R^{ij}(t), \quad \langle \xi^i(t) \xi^j(0) \rangle = G_{sym}^{ij}(t)$$

$$G_R(t) = -i\theta(t) \langle [\mathcal{F}(t), \mathcal{F}(0)] \rangle, \quad G_{sym}(t) = -\frac{i}{2} \langle \{\mathcal{F}(t), \mathcal{F}(0)\} \rangle$$

- The main goal is to use holography in order to evaluate G_{sym} and G_R for the forces of interest in QGP

The local limit

- For $t \gg t_c$ the autocorrelation time of the force

$$\int_0^\infty dt' \gamma(t') \dot{X}(t-t') \rightarrow \eta \dot{X}(t) \quad , \quad \eta = \int_0^\infty dt' \gamma(t')$$

$$G_{sym}(t-t') \rightarrow \kappa \delta(t-t') \quad , \quad \kappa = \int_0^\infty dt G_{sym}(t)$$

$$\dot{P} + \eta_D P = \xi \quad , \quad \eta_D = \frac{\dot{X}}{P} \eta = \frac{\eta}{\gamma M}$$

- In Fourier space

$$\kappa = G_{sym}(\omega = 0) \quad , \quad \eta = - \lim_{\omega \rightarrow 0} \frac{Im G_R(\omega)}{\omega}$$

- The relation between G_R and G_{sym} is ensemble-dependent. For a thermal ensemble

$$G_{sym}(\omega) = \coth\left(\frac{\omega}{2T}\right) Im G_R(\omega) \quad \Rightarrow \quad \kappa = 2T\eta = 2MT\eta_D$$

we recover the *non-relativistic Einstein relation*.

The holographic strategy

- To determine the stochastic motion of heavy quarks we must therefore calculate the force correlator in QCD as

$$e^{iS_{eff}} = \langle e^{i \int X \mathcal{F}} \rangle$$

- We will calculate them using a holographic dual.
 1. We must identify the force operator \mathcal{F} .
 2. We must solve the classical equations to find the average motion.
 3. Calculate the correlators of the force from the boundary on-shell action using the Son-Starinets prescription for the real-time correlators.

The holographic setup

- There is a 5D bulk described by a general 5D black hole with metric (in the string frame)

$$ds^2 = b^2(r) \left[\frac{dr^2}{f(r)} - f(r)dt^2 + d\vec{x}^2 \right]$$

- The boundary is at

$$r \rightarrow 0 \quad , \quad f \rightarrow 1 \quad , \quad b \rightarrow \frac{\ell}{r} + \dots$$


and at the BH horizon

$$r \rightarrow r_h \quad , \quad f(r_h) = 0 \quad , \quad 4\pi T = |\dot{f}(r_h)|$$

- This is the holographic description of a general strongly coupled plasma (deconfined phase) in a heat bath.

Classical Heavy quark motion

- A heavy quark is modeled by a string moving in the BH background with (constant) velocity \vec{v} .



*Herzog+Karch+kovtun+Kozcac+Yaffe, Gubser
Casadelrrey-Solana+Teaney, Liu+Rajagopal+Wiedeman*

- The dynamics of the string is given by the Nambu-Goto action

$$S_{NG} = -\frac{1}{2\pi\ell_s^2} \int d^2\xi \sqrt{\det \hat{g}} \quad , \quad \hat{g}_{\alpha\beta} = g_{\mu\nu} \partial_\alpha X^\mu \partial_\beta X^\nu$$

The drag force and the world-sheet black hole

- The drag (friction) force is directly calculated by solving for the classical string configuration.
- The induced metric on world-sheet is that of a *two dimensional asymptotically AdS₂ black hole*. Its horizon r_s is at $f(r_s) = v^2$.
- It has Hawking temperature T_s that is **different** from that of the heat-bath.
- The thermal ensemble associated to that black hole controls the force correlators.
- **The Fluctuation/Dissipation relation measures temperature = T_s** . In general T_s depends on T, Λ, v .
- In the conformal case, $T_s = \frac{T}{\sqrt{\gamma}}$. $T_s \rightarrow T$ as $v \rightarrow 0$. *Giecold+Iancu+Mueller, 2009*
- **In all examples we analyzed, $T_s \leq T$ but we can not prove it in general.**
- We always have $0 \leq r_s \leq r_h$. $r_s = 0$ when $v = 1$ and $r_s = r_h$ when $v = 0$.

Fluctuations of the trailing string

- So far we have calculated the average *damped* motion of the trailing string.

- To study the fluctuations we set

$$\vec{X}(r, t) = (vt + \xi(r)) \frac{\vec{v}}{v} + \delta\vec{X}(r, t)$$

- From the boundary coupling

$$S_{bdr} = \int dt X_i(t) \mathcal{F}^i(t) \simeq S_{bdr}^0 + \int dt \delta X_i(t) \mathcal{F}^i(t)$$

- Correlators of \mathcal{F} in the dual QFT are given by holographic correlators of $\delta X_i(t)$ in the bulk string theory.
- They can be obtained according to the standard holographic prescriptions by solving the second order fluctuation equations for $\delta X_i(t)$ on the asymptotically AdS string world-sheet.
- They are **different** for longitudinal and transverse fluctuations. They differ by a multiplicative factor Z that depends on the bulk geometry.

The diffusion constants

- From direct calculation of the IR asymptotics of fluctuation correlators we obtain

$$\kappa^\perp = \frac{b^2(r_s)}{\pi \ell_s^2} T_s \quad , \quad \kappa^\parallel = \frac{b^2(r_s)}{\pi \ell_s^2} \frac{(4\pi)^2}{f'(r_s)^2} T_s^3$$

- We also obtain the relation

$$G_{sym}^i(\omega) = \coth\left(\frac{\omega}{2T_s}\right) G_R^i(\omega)$$

- Because the diffusion and friction coefficients are generically momentum dependent there are non-trivial relations between Langevin equations for momenta and position fluctuations.

- We obtain the **modified Einstein relations**

$$\kappa^\perp = 2\gamma M T_s \eta_D = 2E T_s \eta_D \quad , \quad \kappa^\parallel = 2\gamma^3 M T_s \left[\eta_D + \gamma M v \frac{\partial \eta_D}{\partial p} \right]$$

to be compared with the standard one $\kappa = 2MT\eta_D$.

Validity of local Langevin evolution

- The validity of the local approximation demands that

$$t \gg t_{\text{correlation}} \sim \frac{1}{T_s}$$

- For $(\Delta p^\perp)^2$ to be characterized by κ^\perp we must have

$$t \ll t_{\text{relaxation}} \sim \frac{1}{\eta_D}$$

Therefore we need

$$\frac{1}{\eta_D} \ll \frac{1}{T_s}$$

If this fails we need the full non-local (in time) Langevin evolution.

This translates into an upper bound for the momentum ultra-relativistic quarks of the form

$$p \ll \frac{1}{4} \left(\frac{\ell_s}{\ell} \right)^4 \frac{M_q^3}{T^2} \lambda_s^{-8/3}.$$

Calculations in Improved Holographic QCD

- This is Einstein-dilaton gravity with

$$S = M^3 N_c^2 \int d^5x \sqrt{g} \left[R - \frac{4}{3} \frac{\partial \lambda^2}{\lambda^2} - V(\lambda) \right]$$

- λ is approximately the QCD 't Hooft coupling

- $V(\lambda) = \frac{12}{\ell^2} [1 + c_1 \lambda + c_2 \lambda^2 + \dots]$, $\lambda \rightarrow 0$

- $V(\lambda) \sim \lambda^{\frac{4}{3}} \sqrt{\log \lambda}$, $\lambda \rightarrow \infty$

- It has two phenomenological parameters.

- It agrees well with pure YM, both a zero and finite temperature.

Gursoy+Kiritsis+Mazzanti+Nitti, 2007-2009

YM Entropy




Figure 4: (Color online) Same as in fig. 1, but for the s/T^3 ratio, normalized to the SB limit.

From M. Panero, arXiv:0907.3719

Equation of state





Figure 2: (Color online) Same as in fig. 1, but for the Δ/T^4 ratio, normalized to the SB limit of p/T^4 .


From M. Panero, arXiv:0907.3719

Locality of Langevin evolution





The quantity T_s/η_D is plotted against quark momentum, for different bulk temperatures. Figures refer to the charm and bottom quark, respectively. For each temperature, the validity of the local Langevin equation constrains p to the left of the corresponding vertical line, which marks the transition of T_s/η_D across unity.

- In this case we must obtain a force correlator that vanishes fast enough as $t \rightarrow \infty$.
- In order for this to happen, $\lim_{\omega \rightarrow \infty} \rho(\omega) = 0$.
- We must define dressed force correlators by subtracting the $T = 0$ contributions




Breakdown of the dragging string setup




- $T = T_c$: $P^{charm} \ll 1.5$ TeV and $P^{bottom} \ll 56$ TeV.
- $T = 2T_c$: $P^{charm} \ll 290$ GeV and $P^{bottom} \ll 10$ TeV.
- $T = 3T_c$: $P^{charm} \ll 110$ GeV and $P^{bottom} \ll 4$ TeV.

Symmetric Transverse, Langevin Correlator




The symmetric correlator of the \perp modes by the numerical evaluation (solid line) in the $M_q \rightarrow \infty$ limit. We show in each plot the curves corresponding to the velocities $v = 0.1, 0.9, 0.99$ and different plots for the temperatures $T = T_c, 3T_c$.


Jet quenching parameters




\hat{q}_\perp charm



\hat{q}_\parallel charm



\hat{q}_\perp bottom



\hat{q}_\parallel bottom

The quantities \hat{q}_\perp/T_c^3 and \hat{q}_\parallel/T_c^3 plotted as a function of the quark momentum p . The plots for the charm and the bottom quark differ by a scaling of the horizontal direction.

Outlook

- The Langevin diffusion of heavy quarks in the QGP may be an interesting observable that will provide extra clues for the dynamics in the deconfined phase.
- The relativistic Langevin dynamics from holography is providing a novel paradigm with asymmetric evolution and is expected to be valid in QCD.
- The jet quenching transport coefficients may be calculated and provide important input for the evolution of heavy quarks.
- They thermalize at a temperature dictated by a world-sheet black hole and is distinct from the plasma temperature.
- The local Langevin evolution breaks down for the charm rather early and full correlators are needed (for LHC). These are captured by a WKB analysis.
- A detailed simulation done recently (Akamatsu+Hatsuda+Hirano) used a different Langevin evolution that is not in accord with the one derived via holography
- A new simulation seems necessary in order to test (qualitatively at least the holographic templates and predictions).
- The holographic calculations have ample room for improvement, most importantly by including the fundamental degrees of freedom in the plasma.

Thank you for your Patience

Bibliography

- **REVIEW**: U. Gursoy, E. Kiritsis, L. Mazzanti, G. Michalogiorgakis and F. Nitti, *"Improved Holographic QCD"*.
[ArXiv:1006.5461][[hep-th]],
- U. Gursoy, E. Kiritsis, L. Mazzanti and F. Nitti, *"Langevin diffusion of heavy quarks in non-conformal holographic backgrounds"*
[ArXiv:1006.3261][[hep-th]],.
- U. Gursoy, E. Kiritsis, G. Michalogiorgakis and F. Nitti, *"Thermal Transport and Drag Force in Improved Holographic QCD"*
JHEP 0912:056,2009, [ArXiv:0906.1890][[hep-ph]],.
- U. Gursoy, E. Kiritsis, L. Mazzanti and F. Nitti, *"Improved Holographic Yang-Mills at Finite Temperature: Comparison with Data."*
Nucl.Phys.B820:148-177,2009, [ArXiv:0903.2859][[hep-th]],.
- E. Kiritsis, *"Dissecting the string theory dual of QCD."*
Fortsch.Phys.57:396-417,2009, [ArXiv:0901.1772][[hep-th]],.
- U. Gursoy, E. Kiritsis, L. Mazzanti and F. Nitti, *"Thermodynamics of 5D Dilaton-gravity."*
JHEP 0905 (2009) 033; [ArXiv:0812.0792][[hep-th]],.
- U. Gursoy, E. Kiritsis, L. Mazzanti and F. Nitti, *"Deconfinement and Gluon-Plasma Dynamics in Improved Holographic QCD,"*
Phys. Rev. Lett. 101, 181601 (2008) [ArXiv:0804.0899][[hep-th]],.
- U. Gursoy and E. Kiritsis, *"Exploring improved holographic theories for QCD: Part I,"*
JHEP 0802 (2008) 032[ArXiv:0707.1324][[hep-th]].

- U. Gursoy, E. Kiritsis and F. Nitti,
“Exploring improved holographic theories for QCD: Part II,”
JHEP **0802** (2008) 019[ArXiv:0707.1349][[hep-th]].
- Elias Kiritsis and F. Nitti
On massless 4D gravitons from asymptotically AdS(5) space-times.
Nucl.Phys.**B772** (2007) 67-102;[arXiv:hep-th/0611344]
- R. Casero, E. Kiritsis and A. Paredes,
“Chiral symmetry breaking as open string tachyon condensation,”
Nucl. Phys. B **787** (2007) 98;[arXiv:hep-th/0702155].

The large- N_c expansion in QCD

- The generalization of QCD to N_c colors, has an extra parameter: the theory simplifies in a sense when $N_c \rightarrow \infty$.

t 'Hooft 1974

- It has the structure of a string theory, with $g_s \sim \frac{1}{N_c}$. When $N_c = \infty$ the theory contains an infinite number of particles with finite masses and no interactions. The “string” is the “flux tube” of confined color flux that binds quarks and glue together.
- Therefore, at $N_c = \infty$ the theory is “free”.
- The particles are color singlets (glueballs, mesons and baryons).
- It is therefore a good starting point for a perturbative expansion in $\frac{1}{N_c}$.
- There is always the usual coupling constant: $\lambda \equiv a_s N_c$.

- it turns out that $N_c = 3$ is not that far from $N_c = \infty$

Alas, even the leading order in QCD (classical at large N_c) is not easy to compute.

- If $\lambda \ll 1$ we compute in perturbation theory
- This is not the case in QCD at low energy.

AdS/CFT correspondence and holography

- A new twist to the large- N_c expansion was added from standard string theory.

Maldacena 1997

- It involved a cousin theory to QCD: $\mathcal{N}=4$ sYM theory. This is a scale invariant theory: the 't Hooft coupling λ does not run.
- It is claimed to be equivalent to a ten-dimensional (IIB) string theory propagating on a curved space $AdS_5 \times S^5$
- ♠ At strong coupling $\lambda \rightarrow \infty$ the string is stiff, therefore we can approximate it with a point-particle, \rightarrow (super)-gravity approximation.
- we obtain a duality: (a) at $\lambda \rightarrow 0$ perturbative description in terms of gauge theory (b) at $\lambda \rightarrow \infty$ perturbative description in terms of supergravity

Holography in Anti-de-Sitter space

- AdS_5 = maximally symmetric, with negative curvature
- A space with a "radial" direction, where each slice $r = constant$ is a Minkowski₄ space.
- The radial direction can be thought of as an RG scale ($r \sim \frac{1}{E}$): $r=0$ (boundary) is the UV, while $r = \infty$ is the IR.
- It has a single boundary at $r = 0$.
- The gravity fields are "dual" to sYM operators: $g_{\mu\nu} \sim T_{\mu\nu}$, $\phi \sim Tr[F^2]$ etc. One can think of them as "composites".
- The string theory effective action is capturing the dynamics of such "composites"
- Closed strings generate the glueballs. Open strings the mesons. Baryons are more complicated (solitons).

- There have been many non-trivial tests of AdS/CFT correspondence
- The gravity approximation is a (important) bonus because we cannot solve (yet) such string theories.
- But $\mathcal{N} = 4$ sYM is not QCD. How can we describe QCD?
- The problem is the weak coupling in the UV
- ♠ One can add a “phenomenological twist”: write a (gravity) theory that has the features of QCD and is motivated from holography/string theory.
- ♠ The simplest model is known as AdS/QCD: its AdS space with an IR cutoff: its advantage is that is simple. The flip-side is that it has no real dynamics and the coupling does not run.
Polchinski+Strassler 2001, Erlich+Katz+Son+Stephanov 2005, DaRold+Pomarol 2005
- The state of the art: Improved Holographic QCD
Gursoy+Kiritsis+Nitti 2007

Correlators

$$G_R(t) = -i\theta(t)\langle[\mathcal{F}(t), \mathcal{F}(0)]\rangle \quad , \quad G_A(t) = i\theta(-t)\langle[\mathcal{F}(t), \mathcal{F}(0)]\rangle$$


$$G_{sym}(t) = -\frac{i}{2}\langle\{\mathcal{F}(t), \mathcal{F}(0)\}\rangle \quad , \quad G_{anti-sym}(t) = -\frac{i}{2}\langle[\mathcal{F}(t), \mathcal{F}(0)]\rangle$$

$$\langle T\mathcal{F}(t)\mathcal{F}(0)\rangle \equiv \theta(t)\langle\mathcal{F}(t)\mathcal{F}(0)\rangle + \theta(-t)\langle\mathcal{F}(0)\mathcal{F}(t)\rangle = G_{sym} + \frac{1}{2}(G_R + G_A)$$

RETURN

Jet quenching influence

- A non-zero value for the jet quenching parameter for light quarks is essential in explaining the PHIC data. Below we show the nuclear modification factor



Eskola et al. 2005

RETURN

The Kramers Equation

- The Brownian motion induced by the Langevin equation can be remodeled as an evolution in phase space
- Let $P(x^i, p^i, t) d^3x d^3p$ be the probability of an ensemble of probes. The Langevin evolution translates to

$$\left(\frac{\partial}{\partial t} + \frac{\vec{p}}{E} \cdot \frac{\partial}{\partial \vec{x}} \right) P = \frac{\partial}{\partial p^i} \left(\eta_D^{ij} p^j + \frac{1}{2} \kappa^{ij} \frac{\partial}{\partial p^j} \right) P$$

- The equilibrium distribution in a homogeneous ensemble is expected to satisfy,

$$\frac{\partial}{\partial p^i} \left(\eta_D^{ij} p^j + \frac{1}{2} \frac{\partial}{\partial p^j} \kappa^{ij} \right) P = 0$$

- It will be a (non-relativistic) Boltzmann distribution $P \sim e^{-\frac{E}{T}}$ if the Einstein relation holds

$$\kappa^{ij} = 2MT \eta_D^{ij} \quad , \quad E = \frac{\vec{p}^2}{2M}$$

where T is the bath temperature.

Solution of the Langevin Equation

$$\dot{p} = -\eta p + \xi \quad , \quad \langle \xi(t)\xi(t') \rangle = \kappa \delta(t - t')$$

with solution

$$p(t) = p(0)e^{-\eta t} + \int_0^t dt' e^{\eta(t'-t)} \xi(t')$$

$$\langle p(t) \rangle = p(0)e^{-\eta t}$$


$$\langle p(t)^2 \rangle - \langle p(t) \rangle^2 = \int_0^t dt' e^{\eta(t'-t)} \int_0^t dt'' e^{\eta(t''-t)} \langle \xi(t)\xi(t') \rangle = \frac{\kappa}{2\eta} (1 - e^{-2\eta t})$$

- Long times: $t \gg \frac{1}{\eta}$: $\langle p \rangle \rightarrow 0$ and $\langle \Delta p^2 \rangle \rightarrow \frac{\kappa}{2\eta}$.
- Short times: $t \ll \frac{1}{\eta}$: $\langle p \rangle \simeq p(0)$ and $\langle \Delta p^2 \rangle \rightarrow \kappa t$.
- Consider a multidimensional motion and separate

$$\vec{p} = p^{\parallel} + p^{\perp} \quad , \quad \vec{v} \cdot p^{\perp} = 0$$

- The transverse momentum obeys a Langevin process with (by definition) $\langle p^\perp \rangle = 0$ but with an increasing dispersion

$$\langle (\Delta p^\perp)^2 \rangle \rightarrow 2\kappa^\perp t$$



- This defines the “jet quenching parameter”

$$\hat{q} = \frac{\langle (\Delta p^\perp)^2 \rangle}{vt} = 2\frac{\kappa^\perp}{v}$$

- This is a *transport coefficient*.
- Its value is an important ingredient of measured quantities in HIC like R_{AA} .

Schwinger-Keldysh derivation

Consider a system with degrees of freedom $\{ Q \}$ and density matrix $\rho(Q, Q', t)$. that evolves as

$$\rho(Q_f, Q'_f, t) = U(Q_f, Q_0, t, t_0) \rho(Q_0, Q'_0, t_0) U^\dagger(Q'_f, Q'_0, t, t_0)$$


where the evolution operator is given by the path integral

$$U(Q_f, Q_0, t, t_0) = \int DQ e^{iS(Q)} = \int DQ e^{i \int_{t_0}^t L(Q, \dot{Q})} , \quad Q(t_0) = Q_0 \quad , \quad Q(t) = Q_f$$

Therefore the density matrix is a double path integral

$$\rho(Q_f, Q'_f, t) = \int DQ \int DQ' e^{i(S(Q) - S(Q'))} \rho(Q_0, Q'_0, t_0)$$

It is natural to double the fields, call $Q = Q_+$, $Q' = Q_-$, and consider Q_\pm the values of Q on the double (Keldysh) contour



We now consider a single particle described by $X(t)$, and a statistical ensemble, described by a QFT with degrees of freedom $\Phi(x, t)$. We assume a linear interaction between X and some functional $\mathcal{F}(t)$ of the QFT fields Φ .

$$S = S_0(X) + S_{QFT}(\Phi) + S_{int}(X, \Phi) \quad , \quad S_{int}(X, \Phi) = \int dt X(t)\mathcal{F}(t)$$

We assume that the particle starts at $X = x_i$ at $t_i = -\infty$

$$\rho_i = \delta(X - x_i)\delta(X' - x_i)\rho_i(\Phi, \Phi')$$

We would like to compute the reduced density matrix at time t :

$$\rho(X, X', t) = Tr_{\Phi}\rho(X, X', \Phi, \Phi', t)$$

That we can now write as a path integral using a doubled set of fields

$$\rho(X, X', t) = \int DX_+ \int DX_- e^{iS_0(X_+) - iS_0(X_-)} \int D\Phi_+ D\Phi_- e^{iS_+(X_+, \Phi_+) - iS_-(X_-, \Phi_-)} \rho_i(\Phi_+, \Phi_-)$$

where the the trace in the QFT path integral is obtained by setting $(\Phi_+)_f = (\Phi_-)_f$ and

$$S_{\pm} = S_{QFT} + \int X\mathcal{F}$$

Therefore the effective density matrix evolves according to the effective action

$$S_{eff}(X_+, X_-) = S_0(X_+) - S_0(X_-) + S_{IF}(X_+, X_-)$$

$$e^{iS_{IF}} = \langle e^{i \int X_+ \mathcal{F}_+ - i \int X_- \mathcal{F}_-} \rangle_{\text{QFT ensemble}}$$

Feynman+Vernon, 1963

We expand the exponential to quadratic order

$$\begin{aligned}
 \langle e^{i \int X_+ \mathcal{F}_+ - i \int X_- \mathcal{F}_-} \rangle_{\text{QFT ensemble}} &\simeq 1 + i \int dt \langle \mathcal{F}(t) \rangle (X_+ - X_-) - \\
 -\frac{i}{2} \int dt \int dt' &[-X_+(t) i \langle \mathcal{F}_+(t) \mathcal{F}_+(t') \rangle X_+(t') + X_-(t) i \langle \mathcal{F}_-(t) \mathcal{F}_+(t') \rangle X_+(t') + \\
 + X_+(t) i \langle \mathcal{F}_+(t) \mathcal{F}_-(t') \rangle X_-(t') &- X_-(t) i \langle \mathcal{F}_-(t) \mathcal{F}_-(t') \rangle X_-(t')] \\
 &\simeq \exp \left[i \int dt \langle \mathcal{F}(t) \rangle (X_+ - X_-) - \frac{i}{2} \int X_a(t) G_{ab}(t, t') X_b(t') \right]
 \end{aligned}$$

with



$$G_{ab}(t, t') \equiv i \langle \mathcal{P} \mathcal{F}_a(t) \mathcal{F}_b(t') \rangle$$

with \mathcal{P} being path ordering along the keldysh contour:

- $+$ operators are time-ordered, $-$ operators are anti-time-ordered
- $-$ operators are always in the future of $+$ operators.



$$G_{+++}(t, t') = -i \langle T \mathcal{F}_+(t) \mathcal{F}_+(t') \rangle$$

$$G_{-+}(t, t') = -i \langle \mathcal{F}_-(t) \mathcal{F}_+(t') \rangle$$



$$G_{+-}(t, t') = -i \langle \mathcal{F}_-(t') \mathcal{F}_+(t) \rangle$$

$$G_{--}(t, t') = -i \langle \bar{T} \mathcal{F}_-(t) \mathcal{F}_-(t') \rangle$$



The Keldysh propagators can be written in terms of the standard ones:

$$G_R(t) = -i\theta(t)\langle[\mathcal{F}(t), \mathcal{F}(0)]\rangle \quad , \quad G_A(t) = i\theta(-t)\langle[\mathcal{F}(t), \mathcal{F}(0)]\rangle$$

$$G_{sym}(t) = -\frac{i}{2}\langle\{\mathcal{F}(t), \mathcal{F}(0)\}\rangle \quad , \quad G_{anti-sym}(t) = -\frac{i}{2}\langle[\mathcal{F}(t), \mathcal{F}(0)]\rangle$$

$$\langle T\mathcal{F}(t)\mathcal{F}(0)\rangle \equiv \theta(t)\langle\mathcal{F}(t)\mathcal{F}(0)\rangle + \theta(-t)\langle\mathcal{F}(0)\mathcal{F}(t)\rangle = G_{sym} + \frac{1}{2}(G_R + G_A)$$

$$G_{++} = G_{sym} + \frac{1}{2}(G_R + G_A) \quad , \quad G_{--} = G_{sym} - \frac{1}{2}(G_R + G_A)$$

$$G_{+-} = G_{sym} + \frac{1}{2}(-G_R + G_A) \quad , \quad G_{-+} = G_{sym} + \frac{1}{2}(G_R - G_A)$$

$$G_{++} + G_{--} - G_{+-} - G_{-+} = 0$$

Using this we can rewrite the effective action as

$$S_{eff} = S_0(X_+) - S_0(X_-) + \int (X_+ - X_-)G_R(X_+ + X_-) + \frac{1}{2}(X_+ - X_-)G_{sym}(X_+ - X_-) \int$$

We now define

$$X_{class} = \frac{1}{2}(X_+ + X_-) \quad , \quad y = X_+ - X_-$$

In the semiclassical limit $y \ll X_{\text{class}}$ and we can expand

$$S_0(X_+) - S_0(X_-) \simeq \int dt \frac{\delta S_0}{\delta X_{\text{class}}} y + \mathcal{O}(y^3)$$

to obtain

$$S_{\text{eff}} = \int dt y(t) \left[\frac{\delta S_0}{\delta X_{\text{class}}(t)} + \int dt' G_R(t, t') X_{\text{class}}(t') \right] + \frac{1}{2} \int dt \int dt' y(t) G_{\text{sym}}(t, t') y(t')$$

Therefore the X path integral becomes

$$Z = \int DX_{\text{class}} \int Dy e^{i \int dt y(t) \left[\frac{\delta S_0}{\delta X_{\text{class}}(t)} + \int dt' G_R(t, t') X_{\text{class}}(t') \right] + \frac{1}{2} \int dt \int dt' y(t) G_{\text{sym}}(t, t') y(t')}$$

We integrate-in a gaussian variable $\xi(t)$ with variance G_{sym} . This will linearize the y integration

$$Z = \int D\xi \int DX_{\text{class}} \int Dy \exp \left[i \int dt y \left(\frac{\delta S_0}{\delta X_{\text{class}}} + G_R X_{\text{class}} - \xi \right) - \frac{1}{2} \xi G_{\text{sym}} \xi \right]$$

Integrating over y we obtain a δ functional,

$$Z = \int D\xi \int DX_{\text{class}} \delta \left(\frac{\delta S_0}{\delta X_{\text{class}}} + G_R X_{\text{class}} - \xi \right) e^{-\frac{1}{2} \xi G_{\text{sym}} \xi}$$

Therefore the path integral is localized in a solution of the generalized Langevin equation

$$\frac{\delta S_0}{\delta X_{\text{class}}(t)} + \int_{-\infty}^t dt' G_R(t, t') X_{\text{class}}(t') = \xi(t) \quad , \quad \langle \xi(t) \xi(t') \rangle = G_{\text{sym}}(t, t')$$

The drag force

- The classical dragging string solution is

$$X^\perp = 0 \quad , \quad X^\parallel = vt + \xi(r) \quad , \quad \xi(0) = 0$$

$$\xi'(r) = \frac{C}{f(r)} \sqrt{\frac{f(r) - v^2}{b^4(r)f(r) - C^2}} \quad , \quad f(r_s) = v^2 \quad , \quad C = b^2(r_s)f(r_s)$$

- The “drag” force is in the longitudinal direction

$$\frac{dp^\parallel}{dt} = -\frac{b^2(r_s)}{2\pi\ell_s^2} v = -\eta_D^{class} p^\parallel \quad , \quad \eta_D^{class} = \frac{1}{M\gamma} \frac{b^2(r_s)}{2\pi\ell_s^2} \quad , \quad \gamma = \frac{1}{\sqrt{1-v^2}}$$

Gursoy+Kiritsis+Michalogiorgakis+Nitti,2009

RETURN

The world-sheet black hole

- Change coordinates to

$$t = \tau + \zeta(r) \quad , \quad \zeta' = \frac{v\xi'}{f - v^2}$$

and write the induced world-sheet metric as

$$ds^2 = b^2(r) \left[-(f(r) - v^2)d\tau^2 + \frac{b^4(r)}{b^4(r)f(r) - C^2}dr^2 \right] \quad ,$$


- This has a (world-sheet) horizon at $r = r_s$.
- It is an asymptotically AdS_2 , two-dimensional black-hole.
- The Hawking temperature can be calculated to be

$$T_s = \frac{1}{4\pi} \sqrt{f(r_s)f'(r_s) \left[4\frac{b'(r_s)}{b(r_s)} + \frac{f'(r_s)}{f(r_s)} \right]}$$

- In general T_s depends on T, Λ, v . In the conformal case, $T_s = \frac{T}{\sqrt{\gamma}}$.
Giegold+Iancu+Mueller, 2009

$$T_s \rightarrow T \quad \text{as} \quad v \rightarrow 0 \quad , \quad T_s \rightarrow \frac{T}{\sqrt{\gamma}} \quad \text{as} \quad v \rightarrow 1$$

- In all examples we analyzed, $T_s \leq T$.
- We always have $0 \leq r_s \leq r_h$. $r_s = 0$ when $v = 1$ and $r_s = r_h$ when $v = 0$.



RETURN

String fluctuations and force correlators

In the diagonal world-sheet frame

$$S_{NG}^{(2)} = -\frac{1}{2\pi\ell_s^2} \int d\tau dr \frac{H^{\alpha\beta}}{2} \left[\frac{\partial_\alpha X^\parallel \partial_\beta X^\parallel}{Z^2} + \sum_{i=1}^2 \partial_\alpha X_i^\perp \partial_\beta X_i^\perp \right]$$

and the fluctuation equations are

$$\partial_\alpha (H^{\alpha\beta} \partial_\beta) X^\perp = 0 \quad , \quad \partial_\alpha \left(\frac{H^{\alpha\beta}}{Z^2} \partial_\beta \right) X^\parallel = 0$$

$$H^{\alpha\beta} = \begin{pmatrix} -\frac{b^4}{\sqrt{(f-v^2)(b^4f-C^2)}} & 0 \\ 0 & \sqrt{(f-v^2)(b^4f-C^2)} \end{pmatrix} \quad , \quad Z \equiv b^2 \sqrt{\frac{f-v^2}{b^4f-C^2}}$$

We look for harmonic solutions $\delta X(r, t) = e^{i\omega\tau} \delta X(r, \omega)$

$$\partial_r \left[\sqrt{(f-v^2)(b^4f-C^2)} \partial_r (\delta X^\perp) \right] + \frac{\omega^2 b^4}{\sqrt{(f-v^2)(b^4f-C^2)}} \delta X^\perp = 0$$

$$\partial_r \left[\frac{1}{Z^2} \sqrt{(f-v^2)(b^4f-C^2)} \partial_r (\delta X^\parallel) \right] + \frac{\omega^2 b^4}{Z^2 \sqrt{(f-v^2)(b^4f-C^2)}} \delta X^\parallel = 0$$

- Near the boundary the equations is symmetric

$$\Psi'' - \frac{2}{r} \Psi' + \gamma^2 \omega^2 \Psi = 0 \quad , \quad \Psi(r, \omega) \sim C_s(\omega) + C_v(\omega) r^3 + \dots$$

- Near the world-sheet horizon, $r \rightarrow r_s$

$$\Psi'' + \frac{1}{r_s - r} \Psi' + \left(\frac{\omega}{4\pi T_s (r_s - r)} \right)^2 \Psi = 0 \quad , \quad \Psi(r, \omega) \sim C_{out}(\omega) (r_s - r)^{\frac{i\omega}{4\pi T_s}} + C_{in}(\omega) (r_s - r)^{-\frac{i\omega}{4\pi T_s}}$$

- To calculate the retarded correlator we have

$$S = \int dr d\tau \mathcal{H}^{\alpha\beta} \partial_\alpha \Psi \partial_\beta \Psi \quad , \quad \mathcal{H}^{\alpha\beta} = \begin{cases} \frac{H^{\alpha\beta}}{2\pi\ell_s^2} \quad , \quad \perp, \\ \frac{H^{\alpha\beta}}{2\pi\ell_s^2 Z^2} \quad , \quad \parallel, \end{cases}$$

- For the retarded correlator

$$G_R(\omega) = \mathcal{H}^{r\alpha}(r) \Psi^*(r, \omega) \partial_\alpha \Psi(r, \omega) \Big|_{\text{boundary}} \quad , \quad \Psi(0, \omega) = 1 \quad , \quad \Psi(r \rightarrow r_s, \omega) \sim (r_s - r)^{-\frac{i\omega}{4\pi T_s}}$$


- The metric entering the wave equations for fluctuations is 2d BH metric, with temperature T_s . Using the Schwinger-Keldysh formalism we can show that

$$G_{sym}^i(\omega) = \coth\left(\frac{\omega}{2T_s}\right) G_R^i(\omega)$$

and therefore the temperature entering the fluctuation-dissipation relations is T_s .


- This is NOT the thermal equilibrium relation of the plasma.

Asymmetry factor (Z)





The function $\gamma Z(r_s)$ as a function of velocity, ($\gamma \equiv 1/\sqrt{1-v^2}$), computed numerically varying the velocity, at different temperatures. The dashed line represents the conformal limit, in which $\gamma Z = 1$ exactly.

World-Sheet Hawking temperature




The ratio of the world-sheet temperature to the bulk black hole temperature, as a function of velocity, for different values of the bulk temperature. The dashed line indicates the AdS -Schwarzschild curve, $T_s = T/\sqrt{\gamma}$.

Symmetric Longitudinal Langevin Correlator





Retarded correlators for finite mass quarks



The retarded correlator real and imaginary part for finite but large quark mass, calculated numerically. The mass is chosen equal to that of charm.

Symmetric correlator for finite mass quarks



Langevin diffusion constants

$$\kappa = G_{sym}(\omega = 0) = -2T_s \frac{G_R(\omega)}{\omega} \Big|_{\omega=0}$$

$$ImG_r(r, t) = \frac{\mathcal{H}^{rr}}{2i} \Psi^* \overleftrightarrow{\partial} \Psi = J^r(r, t) \quad , \quad \partial_r J^r = 0$$

We can compute $ImG_R(\omega)$, anywhere, and the easiest is at the horizon, $r = r_s$:

$$\Psi = C_h (r_s - r)^{-\frac{i\omega}{4\pi T_s}} + \dots \quad , \quad ImG_R = \frac{\mathcal{H}^{rr}}{4\pi T_s (r_s - r)} \Big|_{r_s} |C_h|^2 \omega$$

- Ψ can be computed exactly as $\omega \rightarrow 0$

$$\Psi \simeq 1 + \omega \int_0^r \mathcal{H}^{rr}(r') dr' \Rightarrow C_h = 1 \Rightarrow \kappa = \frac{\mathcal{H}^{rr}}{2\pi(r_s - r)} \Big|_{r_s} = \frac{1}{\pi \ell_s^2} \begin{cases} b^2(r_s) T_s \quad , \quad \perp, \\ (4\pi)^2 \frac{b^2(r_s)}{f'(r_s)^2} T_s^3 \quad , \quad \parallel, \end{cases}$$

The diffusion constants, II

- From direct calculation of the IR asymptotics of fluctuation correlators we obtain

$$\kappa^\perp = \frac{b^2(r_s)}{\pi \ell_s^2} T_s \quad , \quad \kappa^\parallel = \frac{b^2(r_s)}{\pi \ell_s^2} \frac{(4\pi)^2}{f'(r_s)^2} T_s^3$$

- We also obtain the relation

$$G_{sym}^i(\omega) = \coth\left(\frac{\omega}{2T_s}\right) G_R^i(\omega)$$

and therefore the temperature entering the fluctuation-dissipation relations is T_s .

- Because the diffusion and friction coefficients are generically momentum dependent there are non-trivial relations between Langevin equations for momenta and position fluctuations.

$$\dot{\vec{p}} = -\eta_D^{\parallel} p^{\parallel} \hat{v} - \eta_D^{\perp} p^{\perp} + \vec{\xi}(t)$$

In configuration space (where all of this is calculated)

$$\gamma M \delta \ddot{X}^{\perp} = -\eta^{\perp} \delta \dot{X}^{\perp} + \xi^{\perp} \quad , \quad \gamma^3 M \delta \ddot{X}^{\parallel} = -\eta^{\parallel} \delta \dot{X}^{\parallel} + \xi^{\parallel}$$

$$\eta^{\perp} = \frac{1}{\gamma M} \eta_D^{\perp} \quad , \quad \eta^{\parallel} = \frac{1}{\gamma^3 M} \left[\eta_D^{\parallel} + \gamma M v \frac{\partial \eta_D^{\parallel}}{\partial p} \right]$$

- We have computed holographically

$$\eta^{\parallel, \perp} = \frac{\kappa^{\parallel, \perp}}{2T_s}$$

which lead to the **modified Einstein relations**

$$\kappa^{\perp} = 2\gamma M T_s \eta_D^{\perp} = 2E T_s \eta_D^{\perp} \quad , \quad \kappa^{\parallel} = 2\gamma^3 M T_s \left[\eta_D^{\parallel} + \gamma M v \frac{\partial \eta_D^{\parallel}}{\partial p} \right]$$

to be compared with the standard one $\kappa = 2MT\eta_D$.

- Consistency check:

$$\eta_D^{\parallel} = \eta_D^{\perp} = \frac{b^2(r_s)}{M\gamma(2\pi\ell_s^2)}$$

satisfies both Einstein relations.

- This type of relativistic Langevin evolution is different from what has been described so far in the mathematical physics literature.

Debasch+Mallick+Ribet, 1997

- The diffusion constants satisfy the general inequality (in the deconfined phase)

$$\frac{\kappa_{\parallel}}{\kappa_{\perp}} = \left(\frac{4\pi T_s}{f'(r_s)} \right)^2 = 1 + 4v^2 \frac{b'(r_s)}{f'(r_s)b(r_s)} \geq 1$$

equality is attained at $v = 0$.

- For systems similar to QCD, the WKB approximation valid for large ω seems to be valid down to very low frequencies, providing an analytical control over the Langevin correlators.

RETURN

Langevin friction terms

We have

$$\dot{\vec{p}} = -\eta_D^{\parallel} p^{\parallel} \hat{v} - \eta_D^{\perp} p^{\perp} + \vec{\xi}(t)$$

To connect to the holographic equations we must rewrite them as equations for δX

$$\dot{\vec{X}} = \vec{v} + \delta\dot{\vec{X}} \quad , \quad \vec{p} = \frac{M\dot{\vec{X}}}{\sqrt{1 - \dot{\vec{X}} \cdot \dot{\vec{X}}}} = \gamma M \vec{v} + \delta\vec{p}$$

We expand to first order to obtain the equations for the position fluctuations

$$\gamma M \delta\ddot{X}^{\perp} = -\eta^{\perp} \delta\dot{X}^{\perp} + \xi^{\perp} \quad , \quad \gamma^3 M \delta\ddot{X}^{\parallel} = -\eta^{\parallel} \delta\dot{X}^{\parallel} + \xi^{\parallel}$$

$$\eta^{\perp} = \frac{1}{\gamma M} \eta_D^{\perp} \quad , \quad \eta^{\parallel} = \frac{1}{\gamma^3 M} \left[\eta_D^{\parallel} + \gamma M v \frac{\partial \eta_D^{\parallel}}{\partial p} \right]$$

We have computed holographically

$$\eta^{\parallel, \perp} = \frac{\kappa^{\parallel, \perp}}{2T_s}$$

which lead to the modified Einstein relations

$$\kappa^\perp = 2\gamma MT_s \eta_D^\perp = 2ET_s \eta_D^\perp \quad , \quad \kappa^\parallel = 2\gamma^3 MT_s \left[\eta_D^\parallel + \gamma Mv \frac{\partial \eta_D^\parallel}{\partial p} \right]$$


to be compared with the standard one $\kappa = 2MT\eta_D$.

- Consistency check

$$\eta_D^\parallel = \eta_D^\perp = \frac{b^2(r_s)}{M\gamma(2\pi\ell_s^2)}$$

satisfies both Einstein relations.


Comparison with $N=4$




The ratio of the diffusion coefficients κ_{\perp} and κ_{\parallel} to the corresponding value in the holographic conformal $\mathcal{N} = 4$ theory (with $\lambda_{\mathcal{N}=4} = 5.5$) are plotted as a function of the velocity v (in logarithmic horizontal scale) The results are evaluated at different temperatures $T = T_c, 1.5T_c, 3T_c$ in the deconfined phase of the non-conformal model.

- If we choose $\lambda = 0.5$ instead of $\lambda = 5.5$ in the conformal case then our result agrees with the conformal result within the 10% level, in the range $v > 0.6$ and for $T > 1.5T_c$.


Jet quenching parameters




The jet-quenching parameters \hat{q}_\perp and \hat{q}_\parallel obtained from the diffusion constants κ_\perp and κ_\parallel , normalized to the critical temperature T_c , are plotted as a function of the velocity v (in a logarithmic horizontal scale). The results are evaluated at different temperatures.




\hat{q}_\perp charm



\hat{q}_\parallel charm




\hat{q}_\perp bottom




\hat{q}_\parallel bottom


The quantities \hat{q}_\perp/T_c^3 and \hat{q}_\parallel/T_c^3 plotted as a function of the quark momentum p . The plots for the charm and the bottom quark differ by a scaling of the horizontal direction.




\hat{q}_\perp charm



\hat{q}_\parallel charm



\hat{q}_\perp bottom



\hat{q}_\parallel bottom

The jet-quenching parameters \hat{q}_\perp and \hat{q}_\parallel plotted as a function of T/T_c , for different quark momenta.

Systematic uncertainties and approximations

- Large- N_c limit
- Lack of a first principles string theory dual for YM
- Not included light quark degrees of freedom in plasma. Can be accommodated to leading order by a recalibration of temperature. But this may not be enough.

Bigazzi+Cotrone+Mas+Paredes+Ramallo+Tarrío

- Finite mass corrections (may be relevant for charm)

General considerations

- It has been observed in some systems with complicated dynamics, that when they are gently stirred in contact with a heat bath, they reach equilibrium at a temperature $T_s > T$.
Cugliandolo+Kurchan+Peliti, 1997
- This is what we have shown to happen to all strongly-coupled holographic systems, with the difference that always here $T_s < T$.
- The following question is a hundred years-old and unsettled: “What are the Lorentz transformation properties of temperature?”
- Our analysis suggest that
 1. A heavy quark probe moving in a plasma acts as a moving thermometer.
 2. It measures temperature via the fluctuation-dissipation relation.
 3. The temperature it measures is $T_s(v, T, \dots)$. Its dependence on temperature and velocity is simple in conformal systems ($T_s = T/\sqrt{\gamma}$) but more complicated in non-conformal systems, and depends in particular on the dynamical mass scales.

Detailed plan of the presentation

- Title page 0 minutes
- Collaborators 1 minutes
- Plan of the presentation 2 minutes
- Introduction 3 minutes
- Brownian motion and Langevin dynamics 5 minutes
- The generalized Langevin equation 7 minutes
- The local limit 9 minutes
- The holographic strategy 10 minutes
- The holographic setup 12 minutes
- Classical Heavy Quark Motion 14 minutes
- The drag force and the world-sheet black hole 16 minutes

- Fluctuations of the trailing string 17 minutes
- The diffusion constants 19 minutes
- The validity of the local approximation 21 minutes
- Calculations in Improved Holographic QCD 23 minutes
- The entropy 23 minutes
- The trace 24 minutes
- Locality of Langevin evolution 26 minutes
- Breakdown of the dragging string setup 27 minutes
- Symmetric Transverse, Langevin Correlator 28 minutes
- Jet Quenching Parameters 29 minutes
- Outlook 30 minutes
- Bibliography 30 minutes
- The large N_c expansion in QCD 30 minutes
- AdS/CFT correspondence and holography 30 minutes
- Holography in AdS space 30 minutes
- Jet quenching influence 32 minutes

- The Kramers equation 34 minutes
- Solution of the Langevin equation 38 minutes
- Correlators 40 minutes
- The SK derivation of the Langevin equation 40 minutes
- The drag force 43 minutes
- The world-sheet black hole 48 minutes
- String fluctuations and force correlators 48 minutes
- Asymmetry factor (Z) 49 minutes
- World-sheet Hawking temperature 50 minutes
- Symmetric Longitudinal, Langevin Correlator 51 minutes
- Retarded correlators for finite mass quarks 52 minutes
- Symmetric correlator for finite mass quarks 53 minutes
- Langevin diffusion constants 53 minutes
- The diffusion constants, II 61 minutes
- Langevin friction terms 61 minutes
- Comparison with $N=4$ 62 minutes
- Jet Quenching Parameters 65 minutes
- Systematic uncertainties 66 minutes
- More general considerations 69 minutes