Holography and Hydrodynamics

Yaron Oz (Tel-Aviv University)

SIXTH CRETE REGIONAL MEETING ON STRING THEORY

Outline

Black Holes and Hydrodynamics

Relativistic Hydrodynamics

Fluid/Gravity Correspondence

Quantum Anomalies

Turbulence

Singularities

Outlook

<□> <圖> < => < => < => < => <0 < @</p>

Collaborators and References

With B. Keren-Zur, C. Eling, G. Falkovich, I. Fouxon, I. Itkin, X. Liu, Y. Neiman, M. Rabinovich.

PRL **101** (2008) 261602, JHEP **0903**, 120 (2009), PLB **680**, 496 (2009), JHEP **1002**, 069 (2010), PLB **694**, 261 (2010), , JFM **644**, 465 (2010), JHEP **1006**, 006 (2010), CMP **52**, 43 (2011), JHEP **1103**, 006 (2011), JHEP **1012**, 086 (2010), JHEP **1103**, 023 (2011), JHEP **1102**, 070 (2011), arXiv:1103.1657, arXiv:1106.2683 and to appear.

(ロ) (同) (三) (三) (三) (三) (○) (○)

Fluid Dynamics and Gravity

 The AdS/CFT correspondence relates fluid dynamics to black hole dynamics: hydrodynamic regime of the correspondence.





Heavy-Ion Collision

 The QCD plasma produced at RHIC and LHC seems to exhibit a strong coupling dynamics α_s(T_{RHIC}) ~ O(1).



Heavy-Ion Collision

- Nonperturbative methods that can be used to study real time dynamics are largely unavailable.
- Lattice QCD methods are inherently Euclidean.
- The AdS/CFT correspondence provides a real-time nonperturbative framework.
- We can use the strong coupling properties of N = 4 gauge theory plasma as a reference point for describing the strongly coupled QCD plasma.

(ロ) (同) (三) (三) (三) (三) (○) (○)

The Shear Viscosity to Entropy Density Ratio $\frac{\eta}{s}$

$$egin{aligned} T_{\mu
u} &= \mathcal{T}_{\mu
u}^{ extsf{ldeal}} + \eta \mathcal{T}_{\mu
u}^{ extsf{Viscous}} \ \partial^{\mu} \mathcal{T}_{\mu
u} &= \mathbf{0} \end{aligned}$$

• In Einstein's gravity: $\frac{\eta}{s} = \frac{1}{4\pi}$ (Policastro et.al.:2001).



Elliptic Flow

- Hydrodynamic simulations at low shear viscosity to entropy ratio are consistent with RHIC Data.
- The elliptic flow parameter is the second Fourier coefficient $v_2 = \langle Cos(2\phi) \rangle$ of the azimuthal momentum distribution $dN/d\phi$

$$rac{dN}{d\phi} \sim 1 + 2v_2 \operatorname{Cos}(2\phi)$$

(日) (日) (日) (日) (日) (日) (日) (日)

Elliptic Flow

Luzum, Romatschke:2008



The Hydrodynamic Modes

 The effective degrees of freedom are charge densities ρ(x, t), and the hydrodynamics equations are conservation laws

$$\partial_t \rho + \partial_i j^i = 0 \tag{1}$$

The constitutive relations express jⁱ in terms of ρ and its derivatives. For instance jⁱ = −D∂ⁱρ, and we get

$$\partial_t \rho - D \partial_i \partial^i \rho = 0 \tag{2}$$

• Writing $\rho(\vec{k},t) = \int d^3x e^{-i\vec{k}\cdot\vec{x}\rho(\vec{x},t)}$ we have

$$\rho(\vec{k},t) = e^{-Dk^2 t} \rho(\vec{k},t=0)$$
(3)

This is the characteristic behaviour of hydrodynamic mode. It has a life $\tau(k) = \frac{1}{Dk^2}$ which is infinite in the limit $k \to 0$.

Relativistic Hydrodybnamics

 Hydrodynamics applies under the condition that the correlation length of the fluid *l_{cor}* is much smaller than the characteristic scale *L* of variations of the macroscopic fields

$$Kn \equiv I_{cor}/L \ll 1 \tag{4}$$

◆□▶ ◆□▶ ◆□▶ ◆□▶ □ のQ@

Hydrodynamics equations are conservation laws

$$\partial_{\mu}T^{\mu\nu} = 0, \qquad \partial_{\mu}J^{\mu}_{a} = 0$$
 (5)

Relativistic Hydrodynamics

- The equations of relativistic hydrodynamics are determined by the constitutive relation expressing $T^{\mu\nu}$ and J^{μ}_{a} in terms of the energy density $\epsilon(x)$, the pressure p(x), the charge densities $\rho_{a}(x)$ and the four-velocity field $u^{\mu}(x)$ satisfying $u_{\mu}u^{\mu} = -1$.
- The constitutive relation has the form of a series in the small parameter $Kn \ll 1$,

$$T^{\mu\nu}(\mathbf{x}) = \sum_{l=0}^{\infty} T^{\mu\nu}_{(l)}(\mathbf{x}), \ T^{\mu\nu}_{(l)} \sim (Kn)^{l}$$
 (6)

(日) (日) (日) (日) (日) (日) (日) (日)

Ideal Hydrodybnamics

 Keeping only the first term in the series gives ideal hydrodynamics and the stress-energy tensor reads

$$T^{\mu\nu}_{(0)} = \epsilon u^{\mu} u^{\nu} + p \left(\eta^{\mu\nu} + u^{\mu} u^{\nu} \right)$$
(7)

The equation of state $\epsilon(p)$ is an additional input.

CFT hydrodynamics: T^μ_μ = 0, ε = 3p ~ T⁴ and the stress-energy tensor reads

$$T^{\mu\nu}_{(0)} = T^4 [\eta^{\mu\nu} + 4u^{\mu}u^{\nu}]$$
(8)

<日 > < 同 > < 目 > < 目 > < 目 > < 目 > < 0 < 0</p>

Viscous Hydrodynamics

• The dissipative hydrodynamics is obtained by keeping I = 1 term in the series. The stress-energy tensor reads (Landau Frame: $u_{\mu}T_{(1)}^{\mu\nu} = 0$)

$$T^{\mu\nu}_{(1)} = -\eta \sigma^{\mu\nu} - \xi (\partial_{\alpha} u^{\alpha}) \left(\eta^{\mu\nu} + u^{\mu} u^{\nu} \right)$$
(9)

where

$$\sigma^{\mu\nu} = (\partial^{\mu}u^{\nu} + \partial^{\nu}u^{\mu} + u^{\nu}u^{\rho}\partial_{\rho}u^{\mu} + u^{\mu}u^{\rho}\partial_{\rho}u^{\nu} - \frac{2}{3}\partial_{\alpha}u^{\alpha}[\eta^{\mu\nu} + u^{\mu}u^{\nu}]$$
(10)

• The dissipative hydrodynamics of a CFT is determined by only one kinetic coefficient - the shear viscosity η

$$T^{\mu\nu}_{(1)} = -\eta \sigma^{\mu\nu}$$
 (11)

<日 > < 同 > < 目 > < 目 > < 目 > < 目 > < 0 < 0</p>

Gravitational Dual Description

 Consider the five-dimensional Einstein equations with negative cosmological constant

$$R_{mn} + 4g_{mn} = 0, \quad R = -20$$
 (12)

 These equations have a particular "thermal equilibrium" solution - the boosted black brane

$$ds^{2} = -2u_{\mu}dx^{\mu}dr - r^{2}f[br]u_{\mu}u_{\nu}dx^{\mu}dx^{\nu} + r^{2}P_{\mu\nu}dx^{\mu}dx^{\nu}$$
(13)

where

$$f(r) = 1 - \frac{1}{r^4}, \ P^{\mu\nu} = u^{\mu}u^{\nu} + \eta^{\mu\nu}$$

and the constant $T = 1/\pi b$ is the temperature.

◆□ ▶ ◆□ ▶ ◆ □ ▶ ◆ □ ▶ ● □ ● ● ● ●

Gravitational Description

 One looks for a solution of the Einstein equation by the method of variation of constants (Bhattacharya et.al.: 2007)

$$g_{mn} = (g_0)_{mn} + \delta g_{mn} \tag{14}$$

$$(g_0)_{mn} dy^m dy^n = -2u_\mu(x^\alpha) dx^\mu dr - r^2 f[b(x^\alpha)r]u_\mu(x^\alpha)u_\nu(x^\alpha)dx^\mu dx^\nu + r^2 P_{\mu\nu}(x^\alpha)dx^\mu dx^\nu$$
(15)

y = (x^μ, r). As in the Boltzmann equation, the condition of constructibility of the series solution produces equations for u^μ(x^α) and T(x^α) = 1/πb(x^α). The series for g_{mn} is the series in the Knudsen number of the boundary CFT hydrodynamics.

Horizon Dynamics

- The way the black brane horizon geometry encodes the boundary fluid dynamics is reminiscent of the *Membrane Paradigm* (Damour) in classical general relativity, according to which any black hole has a fictitious fluid living on its horizon.
- The dynamics of the event horizon of a black brane in asymptotically AdS space-time (Gauss-Codazzi equations) is described by the Navier-Stokes equations.
- Recently the two approaches are related by an RG flow (Bredberg et. al: 2010).
- The apparent horizon emerges as a useful object that captures time dependent properties of the plasma.

The Horizon Geometry

- Thermalization in field theory is the process of black hole creation in gravity.
- Hydrodynamics is the deformation of the black hole geometry.

Uμ(x) T(x)

◆□▶ ◆□▶ ◆□▶ ◆□▶ □ のQ@

The Bulk Viscosity ζ

- (Eling,Y.O.:2011) The bulk viscosity is captured by the horizon dynamics.
- Consider (d + 1)-dimensional gravitational backgrounds holographically describing thermal states in strongly coupled d-dimensional field theories. The (d + 1)-dimensional gravitational action reads

$$I = \frac{1}{16\pi} \int \sqrt{-g} d^{d+1} x \left(\mathcal{R} - \frac{1}{2} \sum_{i} (\partial \phi_i)^2 - V(\phi_i) \right) + I_{gauge} ,$$
(16)

 $V(\phi_i)$ represents the potential for the scalar fields and I_{gauge} represents the action of gauge fields (abelian or non-abelian) A^a_{μ} .

The Bulk Viscosity ζ

- The null horizon focusing equation obtained by projecting the field equations of (16) on the horizon is equivalent via the fluid/gravity correspondence to the entropy balance law of the fluid.
- Using this equation we derived

$$\frac{\zeta}{\eta} = \sum_{i} \left(s \frac{d\phi_{i}^{H}}{ds} + \rho^{a} \frac{d\phi_{i}^{H}}{d\rho^{a}} \right)^{2}$$

 η is the shear viscosity, *s* is the entropy density, ρ^a are the charges associated with the gauge fields A^a_{μ} , and ϕ^H_i are the values of the scalar fields on the horizon.

• $\frac{\zeta}{\eta}$ rises in the vicinity of T_c .

- We denote the coordinates of the bulk spacetime by x^A = (r, x^μ), A = 0, ..., d. The x^μ are local coordinates on the horizon H; r is a transverse coordinate, with r = 0 on H.
- ∂_Ar is a null covector tangent to the *H*. When raised with the bulk metric, it gives a vector field ℓ^A

$$\ell^{A} = g^{AB} \partial_{B} r = (0, \ell^{\mu}) \tag{17}$$

• These choices fix the following components of the inverse bulk metric on \mathcal{H} :

$$g^{rr} = 0; \quad g^{r\mu} = \ell^{\mu}$$
 (18)

(ロ) (同) (三) (三) (三) (三) (○) (○)

- The pullback of g_{AB} into H is the degenerate horizon metric γ_{μν}, γ_{μν}ℓ^ν = 0.
- The Lie derivative of γ_{µν} along ℓ^µ gives us the shear/expansion tensor, or "second fundamental form":

$$\theta_{\mu\nu} \equiv \frac{1}{2} \mathcal{L}_{\ell} \gamma_{\mu\nu} \tag{19}$$

• We can write a decomposition of $\theta_{\mu\nu}$ into a shear tensor $\sigma_{\mu\nu}^{(H)}$ and an expansion coefficient θ :

$$\theta_{\mu\nu} = \sigma_{\mu\nu}^{(H)} + \frac{1}{d-1} \theta \gamma_{\mu\nu}$$
(20)

・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・

Extrinsic Curvature

$$\Theta_{\mu}{}^{\nu} = \nabla_{\mu}\ell^{\nu} \tag{21}$$

▲□▶▲□▶▲□▶▲□▶ □ のQで



(

The extrinsic curvature measures the bending of the surface embedded in the higher dimensional space. The normal vector n(x) at the point x is parallel transported to the normal vector n(y) at the point y and differs from the dashed vector at point y.

• The expansion θ can be expressed as

$$\theta = \mathbf{v}^{-1} \partial_{\mu} (\mathbf{v} \ell^{\mu}) = \mathbf{v}^{-1} \partial_{\mu} \mathbf{S}^{\mu} , \qquad (22)$$

◆□▶ ◆□▶ ◆□▶ ◆□▶ □ のQで

where v is a scalar density equal to the horizon area density, and $S^{\mu} = v I^{\mu}$ is the horizon area current.



The field equations read

$$\mathcal{R}_{AB} - \frac{1}{2}g_{AB}\mathcal{R} = \frac{1}{2}\sum_{i}\left(\partial_{A}\phi_{i}\partial_{B}\phi_{i} + \frac{1}{2}g_{AB}V(\phi_{i})\right) + T_{AB}^{gauge}$$
(23)

 In (radially shifted) Eddington-Finkelstein coordinates the metric solutions that we are interested in take the form

$$g_{AB}dx^{A}dx^{B} = ds_{(0)}^{2} = -c_{T}^{2}(r+R)\ell_{\mu}\ell_{\nu}dx^{\mu}dx^{\nu} + 2c_{R}(r+R)\ell_{\mu}dx^{\mu}dr + c_{X}^{2}(r+R)P_{\mu\nu}dx^{\mu}dx^{\nu}$$
(24)

where at r = 0, the function c_T vanishes and there is a horizon. $P_{\mu\nu} = \eta_{\mu\nu} + \ell_{\mu}\ell_{\nu}$ is the projection tensor.

 The Bekenstein-Hawking entropy density and the Hawking temperature associated with the metric are

$$s = \frac{1}{4}c_X(R)^{d-1}; \quad T = \frac{c_T(R)c'_T(R)}{c_R(R)}$$
 (25)

 In addition there are the solutions to the scalar fields and gauge fields

$$\phi_i(r+R), \qquad A_B^a = A^a(r)\delta_B^t \tag{26}$$

From the solution for the gauge field, one can read off the values of the chemical potentials μ_a and the charge densities ρ^a .

 Projection of the field equations on l^Al^B give the Null Focusing Equation. It can be written as

$$\partial_{\mu}(s\ell^{\mu}) = \frac{1}{4}\partial_{\mu}S^{\mu} = \frac{s}{2\pi T}\pi_{\mu\nu}\pi^{\mu\nu} + \frac{s}{4\pi T}\sum_{i}(D\phi_{i})^{2} + \cdots$$
(27)

where the dots indicate the gauge field terms, and

$$\pi_{\mu\nu} = P^{\rho}_{\mu} P^{\sigma}_{\nu} \partial_{(\rho} \ell_{\sigma)} - \frac{1}{d-1} P_{\mu\nu} \partial_{\rho} \ell^{\rho}$$
(28)

<日 > < 同 > < 目 > < 目 > < 目 > < 目 > < 0 < 0</p>

• We can write

$$D\phi^{H} = \frac{d\phi_{i}^{H}}{ds}Ds + \frac{d\phi_{i}^{H}}{d\rho^{a}}D\rho^{a} = -\left(s\frac{d\phi_{i}^{H}}{ds} + \rho^{a}\frac{d\phi_{i}^{H}}{d\rho^{a}}\right)(\partial_{\rho}\ell^{\rho})$$
(29)

 In the last equality we used the ideal conservation law, and also the ideal conservation equation for any charge current present

$$\partial_{\mu}J^{\mu a} = \partial_{\mu}(\rho^{a}\ell^{\mu}) = 0 \tag{30}$$

 Therefore, the Raychaudhuri equation to second order takes in the fluid/gravity correspondence the form of a fluid entropy balance law

$$\partial_{\mu}(\mathbf{s}\ell^{\mu}) = \frac{2\eta}{T} \pi_{\mu\nu} \pi^{\mu\nu} + \frac{\zeta}{T} (\partial_{\rho}\ell^{\rho})^2$$
(31)

with the standard shear viscosity obeying $\eta/s = 1/4\pi$, but now with an additional bulk viscosity term.

• The formula seems to hold exactly in some cases (Buchel et. al.:2011).

◆□▶ ◆□▶ ◆□▶ ◆□▶ □ のQ@

• In fact it is an exact formula (Eling and Oz: to appear).

Anomalies

 The hydrodynamics description exhibits an interesting effect when a global symmetry current of the microscopic theory is anomalous

$$D_{\mu}J^{\mu}_{lpha}=rac{1}{8}m{C}_{lphaeta\gamma}\epsilon^{\mu
u
ho\sigma}m{F}^{eta}_{\mu
u}m{F}^{\gamma}_{
ho\sigma}$$

 The form of an anomalous symmetry current is modified in the hydrodynamic description by a term proportional to the vorticity of the fluid

$$\omega^{\mu} \equiv \frac{1}{2} \epsilon^{\mu\nu\lambda\rho} u_{\nu} \partial_{\lambda} u_{\rho}$$

 This has been first discovered in the context of the the fluid/gravity correspondence (Erdmenger et.al, Banerjee et.al.:2008).

Anomalies

The global symmetry current takes the form

$$j_{a}^{\mu} = \rho_{a} u^{\mu} + \sigma_{a}^{\ b} \left(E_{b}^{\mu} - T P^{\mu\nu} D_{\nu} \frac{\mu_{b}}{T} \right) + \xi_{a} \omega^{\mu} + \xi_{ab}^{(B)} B^{b\mu}$$

where ρ_a , T, μ_a and σ_a^b are the charge densities, temperature, chemical potentials and the conductivities of the medium.

◆□▶ ◆□▶ ◆□▶ ◆□▶ □ のQ@

Anomalies

The anomaly coefficients are (Son, Surowka; Neiman, Y.O)

$$\begin{aligned} \xi_{a} &= C_{abc}\mu^{b}\mu^{c} + 2\beta_{a}T^{2} - \frac{2n_{a}}{\epsilon + p} \left(\frac{1}{3}C_{bcd}\mu^{b}\mu^{c}\mu^{d} + 2\beta_{b}\mu^{b}T^{2}\right) \\ \xi_{ab}^{(B)} &= C_{abc}\mu^{c} - \frac{n_{a}}{\epsilon + p} \left(\frac{1}{2}C_{bcd}\mu^{c}\mu^{d} + \beta_{b}T^{2}\right) \end{aligned}$$

(日) (日) (日) (日) (日) (日) (日) (日)

 C_{abc} is the coefficient of the triangle anomaly of the currents j_a^{μ}, j_b^{μ} and j_c^{μ} .

• β_a seem to correspond to the T^2J gravitational anomaly (free fermions calculation by Landsteiner et.al: 2011).

- In very energetic collisions the hot dense QCD matter can go through a phase transition to a deconfined phase described by a fluid-like collective motion of quarks and gluons. We consider a deconfined QCD fluid phase, with three light flavors and chiral symmetry restoration.
- We consider the experimental implications of the axial current triangle diagram anomaly in a hydrodynamic description of high density QCD.

(ロ) (同) (三) (三) (三) (三) (○) (○)

Chiral Magnetic and Vortical Effects

• (Kharzeev, Son :2011) Chiral magnetic effect: charge separation. Chiral vortical effect: baryon number separation.

$$ec{J} = rac{N_c \mu_5}{2\pi^2} \left(tr(VAQ) ec{B} + tr(VAB) 2 \mu_B ec{\omega}
ight)$$

The electromagnetic current corresponds to V = Q and the baryon current to V = B.

(ロ) (同) (三) (三) (三) (三) (○) (○)

• The ratio between the baryon asymmetry and the charge asymmetry increases when the center of mass energy is lowered.

- (Keren-Zur,Y.O.:2010): The basic idea is that the the axial charge density, in a locally uniform flow of massless fermions, is a measure of the alignment between the fermion spins.
- When the QCD fluid freezes out and the quarks bind to form hadrons, aligned spins result in spin-excited hadrons. The ratio between spin-excited and low spin hadron production and its angular distribution may therefore be used as a measurement of the axial charge distribution.
- We predict the qualitative angular distribution and centrality dependence of the axial charge.

(ロ) (同) (三) (三) (三) (三) (○) (○)



◆□> <圖> < E> < E> E のQQ

• Our main proposal is that for off-central collisions we expect enhancement of Ω^- production along the rotation axis of the collision





Superfluid Hydrodynamics

- The most remarkable property of liquid helium below the λ-point is superfluidity. It is the ability of the fluid to flow inside narrow capillaries without friction, discovered by Kapitza. The theoretical basis for understanding the phenomenon of superfluidity was given by Landau.
- The hydrodynamics of a superfluid consists of two motions: the motion of the normal part of the fluid, and the motion of the superfluid part which is an irrotational one, i.e. its velocity is curl free.
- A superfluid can be described as a fluid with a spontaneously broken symmetry, where the superfluid component is the condensate, and its velocity is proportional to the Goldstone phase gradient.
- The hydrodynamics of relativistic superfluids is relevant to the study of neutron stars, and highly dense quark matter at the low temperature Color-Flavor locked phase.

Superfluid Hydrodynamics - CFL Phase

• Anomalous transport (Bhattacharya et. al.:2011).



(Neiman, Y.O.) A Chiral Electric Effect:

$$J^{a\mu} = C^{cgd} \left(\delta^{a}_{c} - \frac{n^{a} \mu_{c}}{\epsilon + p} \right) \left(\delta^{b}_{d} - \frac{n^{b} \mu_{d}}{\epsilon + p} \right) \epsilon^{\mu\nu\rho\sigma} u_{\nu} \xi_{g\rho} \left(E_{b\sigma} - T \partial_{\sigma} \frac{\mu_{b}}{T} \right)$$

◆□▶ ◆□▶ ◆□▶ ◆□▶ □ のQで

Relativistic Turbulence

• The Reynolds number is

$$\mathcal{R}_{e} \sim rac{TL}{\eta/s}$$
 (32)

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

When \mathcal{R}_e is large we expect turbulence.

Is there a universal structure in relativistic turbulence?

Relativistic Turbulence

 In relativistic hydrodynamics we consider the hydrodynamics equation with a random force term

$$\partial^{\nu} T_{\mu\nu} = f_{\mu} \tag{33}$$

and derive the exact scaling relation (Fouxon, Y.O.:2010)

 $\langle T_{0j}(0,t)T_{ij}(r,t)\rangle = \epsilon r_i$

where $d\langle T_{0j}(0,t)f_j(0,t)\rangle \equiv \epsilon$.

 In charged hydrodynamics with a conserved symmetry current J^μ:

 $\langle J_0(0,t)J_i(r,t)\rangle = \epsilon r_i$

・ロト・西・・日・・日・・日・

RHIC and LHC

- Does turbulence show up in RHIC and LHC?
- For gold collisions at RHIC, the characteristic scale *L* is the radius of a gold nucleus $L \sim 6$ Fermi, the temperature is the QCD scale $T \sim 200$ MeV, and $\frac{\eta}{s} \sim \frac{1}{4\pi}$ is a characteristic value of strongly coupled gauge theories.
- With these *R_e* is too small for an experimental realization of the steady state relativistic turbulence.

(ロ) (同) (三) (三) (三) (三) (○) (○)

Singularities in Hydrodynamics

- (Y.O.,M.Rabinovich:2010) The basic question concerning singularities in the hydrodynamic description, is whether starting with appropriate initial conditions, where the velocity vector field and its derivatives are bounded, can the system evolve such that it will exhibit within a finite time a blowup of the derivatives of the vector field.
- Physically, such singularities if present, indicate a breakdown of the effective hydrodynamic description at long distances and imply that some new degrees of freedom are required.

(ロ) (同) (三) (三) (三) (三) (○) (○)

Shock Wave



Singularities in Gravity

 The issue of hydrodynamic singularities has an analogue in gravity. Given an appropriate Cauchy data, will the evolving space-time geometry exhibit a naked singularity, i.e. a blowup of curvature invariants and the energy density of matter fields at a point not covered by a horizon.

(ロ) (同) (三) (三) (三) (三) (○) (○)

• Initial conditions violating Penrose inequality lead to a creation of singularities.

Singularities in Gravity

 (I.Itkin,Y.O.:2011) The general form of the Penrose inequality for asymptotically locally AdS_{d+1} spaces, with electric charge q and a boundary topology characterized by k = 0, ±1:

$$M - M_{0} \geq \frac{(d-1)\Omega_{d-1,k}}{16\pi} [q^{2} (\frac{\Omega_{d-1,k}}{A})^{\frac{d-2}{d-1}} + k(\frac{A}{\Omega_{d-1,k}})^{\frac{d-2}{d-1}} + \frac{1}{l^{2}} (\frac{A}{\Omega_{d-1,k}})^{\frac{d}{d-1}}]$$
(34)

where

$$M_0 = \frac{1}{8\pi} (-k)^{d/2} \frac{(d-1)!!^2}{d!} I^{d-2} \Omega_{d-1,k}$$
(35)

for *d* even and zero for *d* odd.

Outlook

- Turbulence and its universal structure.
- Singularities in hydrodynamics and cosmic censorship.
- Anomalous transport in normal fluids and superfluids.
- Applications to heavy ion collisions, QCD at high densities and Neutron stars.



◆□▶ ◆□▶ ◆□▶ ◆□▶ □ のQで